# WATER, LIFE AND MAN

GB652 Fur 1979

... FURTADO



# WATER, LIFE AND MAN

an inaugural lecture delivered at the University of Malaya on Friday, September 8, 1979

by

JOSE I. DOS R. FURTADO BSc(WAust), PhD Professor of Zoology University of Malaya

Kuala Lumpur University of Malaya 1979



# WATER, LIFE AND MAN

"WATER, LIFE AND MAN" are distinct and yet interrelated phenomena which I wish to discuss this evening. I have chosen this title as representative of the umbrella encompassing my research and academic interests and endeavours; and as the umbrella of academic policies I envisage for the advancement of zoology in this University in particular, and in Malaysia and South East Asia in general. I will firstly elaborate on the interactions between water and life that preoccupy me as a zoologist, and secondly discuss the interrelationships between water, life and man that concern me as an ecologist and humanist besides that of a zoologist. What I have to say on "WATER, LIFE AND MAN" has been crystallized by the poet, William Wordsworth in the following lines:

"There was a time when meadow, grove and stream, The earth, and every common sight, To me did seem apparelled in celestial light, The glory and the freshness of a dream.

It is not now as it hath been of yore – Turn wheresoever I may By night or day, The things which I have seen I now can see no more"

- William Wordsworth.

#### Water

Water is the most important fluid on Earth. It is a neutral liquid with supreme solvent properties, large heat capacity, high surface tension, and an anomalous behaviour. Water is a necessary condition for organic

life forming 60-70% of most organisms. It is therefore essential for sustaining human life. However, in spite of its importance, only 5% of water is in a free state and cycles on Earth, while 95% is chemically bound in rocks (Clapham 1973):

(A)	Chemically bound in rocks (i.e. does not cycle)			Amount (x10 <sup>17</sup> kg)
	(1) Crystalline rocks		250,000	
	(2)	Sedimentary rocks		2,100
		Sub-to	tal	252,100 (95%)
(B)	<i>Free water</i> (100%) (i.e. moves via hydrological cycle) (3) Oceans (97.3%)			13,200
	(4)	(4) Icecaps & glaciers (2,1%)		292
	(5)	i) Ground water 4,000 m. (0.6%)		83.5
	(6)	) Freshwater lakes (0.01%)		1.25
	(7)	Saline lakes & inland seas (0.01%)		1.04
	(8)	8) Soil moisture (0.005%)		0.67
	(9)	(9) Atmospheric water vapour (0.001%)		0.13
	(10)	Rivers (0.0001%)		0.013
			Sub-total	13,578.603 (15%)
			Total	265,678.603

Of the 5% free water, 97.3% occurs in a saline form in the oceans, whereas 2.7% occurs in a fresh form on land. Of the total fresh water, 77.1% occurs as ice-sheet and glaciers, 22.1% as ground water, 0.3% as freshwater in lakes, 0.3% as water in saline lakes and inland seas, 0.2% as soil moisture, 0.03% as atmospheric water vapour and 0.003% as water in rivers.

In view of its low stock in the atmosphere in relation to annual precipitation on Earth, fresh water cycles very rapidly through the biosphere at the rate of at least 32 times/year, or at the most 11.4 days/cycle. This rapid cycling is enhanced by the uneven nature of precipitation over land, which is maximum near the equator. This cycling is possible through three pathways of fresh water transfer: (1) evapotranspiration constituting about 70% of the flow, (2) surface runoff about 27% of the flow, and (3) ground water constituting about 3% of the flow.

Transpiration by plants is related to the mechanism of nutrient uptake by the roots, and forms at least 50% of evapotranspiration. It is higher in aquatic plants such as rice which are important in freshwater swamps. Surface runoff is counterbalanced by differential rates of evaporation and precipitation over land and sea, with more evaporation occurring over the sea and more precipitation over the land, and by the movement of water-bearing clouds from sea to the land. It also facilitates erosion and the movement of nutrients downstream. The latter is counterbalanced by weathering processes in a biogeochemical cycle. Ground water flows are extremely variable, being usually slow and resulting in water storage. Its storage capcity is 60 times that of freshwater lakes.

In view of these three pathways, water retention on land varies from 10-to 100 days, exceeding 100 days only in the case of ground water flows. The natural flow of water through these pathways is a very important natural resource. My interest has focussed particularly on surface runoff by rivers and swamps in South East Asia, which is dependent on the nature of evapotranspiration and ground water flows on the one hand, and on the nature of monsoonal winds and thermal instability governing rainfall, on the other. On the basis of these forces, at least three and up to five hydrological regions can be distinguished in Peninsular Malaysia (Chia 1974; Dale 1959; Stewart 1930; Wycherley 1967). The drainage basins in these regions are characterised by two peak, one-peak or continuous flows per year on the basis of different monsoonal and/or intermonsoonal impacts.

Streams, swamps and rivers form the main drainage system of tropical South East Asia. Except for some volcanic lakes of limited drainage area, there are no natural lakes of major importance to the drainage of this region. There are however some sunken river basins, with the characteristics of both lakes and rivers, such as the great Lake or Tonle Sap in Kampuchea. The drainage system in this tropical region is similar to others on this planet, in that streams and rivers are connected to swamps and flood plains, and that the system is adapted for floods or peak flows. They differ at least in one respect, however, in that the swamps and flood plains in this region have been transformed largely for wet rice cultivation.

The drainage waters of tropical South East Asia are intimately connected to their landscape on a catchment basis, and are dependent on the vegetation, lithology and pedology of the river basin for their nutrient and organic inflows. They are generally acidic, poorly oxygenated, low in biological and chemical oxygen demand, low in suspended solids except during thunderstorms and heavy rains, rich in silicates, and poor in nutrients especially sulphates, calcium and magnesium, because of low nutrient inflows from the landscape and slow mineralization of decomposing organic matter (Johnson 1967a and 1973; Bishop 1971 and 1973; Ho and Furtado 1974; Lim 1974; Cheng 1977; Kobayashi 1959; Mizuno and Mori 1970). These waters may be clear or black in colour depending on the nature of the soils, with the black (or "aver hitam") colour due to podsols or peat soils. These waters are thus similar to those of other tropical drainage basins (Sioli 1975a & b; Fittkau, Irmler, Junk, Reiss and Schmidt 1975). Nevertheless, very few drainage waters have been studied (e.g. Sungei Gombak, Sungei Langat, Sungei Renggam and some streams in South Malaysia) for a comprehensive understanding of their dynamics in tropical South East Asia, leaving much scope for critically needed research.

In forested water catchments interflows are more important than surface runoff in contributing water, organic matter and nutrients to the drainage system, with a contribution of about 80% (Gounaratnam 1974; Manokaran 1974; Sandhu 1978). This denudation is fairly stable. However, when the forests are manipulated for economic production, surface runoff is the dominant source of these materials with a contribution of about 50%, because of lower rainfall interception by the vegetation. These manipulations critically interrupt the different rates of material cycling for essential nutrients such as nitrogen, phosphorus, potassium and calcium (Golley, McGinnis, Clements, Child and Duever

1975) resulting in changes to the structure and function of both the land and drainage system. It results not only in enrichment of the drainage system with nutrients and organic matter (Ho 1973; Bishop 1971 and 1973) but also in enhancing floods, soil erosion, sediment load and turbidity (Pereira 1962; Ho 1973; Leigh and Low 1973; Low 1972: Leigh 1974; Gunaratnam 1974; Sandhu 1978). It has been estimated that suspended sediment load in the drainage system increases by at least one order of magnitude, from 50-100 tons/sq. mile/year to 300-1000 tons/sq. mile/year, when land is transformed from forest to agriculture (Gunaratnam 1974). The intensity of soil erosion and suspended sediment load depend on the pedology, nature of the storm and slope of the land (Leigh 1974; Leigh and Low 1973). Enrichment of the drainage system, flooding, soil erosion, sediment load and turbidity may be regulated in various ways (Leigh and Low 1973), especially by the use of perennial tree plantation crops (Swank and Douglas 1974) and by increasing the density of trees (Teoh 1974). The landscape vegetation in gross units (i.e. forests, agricultural plantations, and urban areas) is related to water quantity and quality in the drainage system (Sandhu 1978; Sandhu, Gunaratnam & Furtado, in press). However, there is a need to define these relationships in terms of specific plant communities for predictive purposes, using modelling technique.

Although organic matter transfer from the landscape to the drainage system determines the structure and function of rivers, there are no previous studies in tropical South East Asia on the decomposition and fate of this matter, and mineralization processes in the system (Verghese 1975; Verghese and Furtado, in press).

## Life and Water

Life is a very special phenomenon. It is a behavioural pattern which matter exhibits when it reaches certain levels of organization and complexity. Living organisms first appeared in the deeper waters of tropical seas about 3.5 billion years ago, where they were protected against the effects of radiation. These early life forms were essentially heterotrophic, dependent on high energy compounds for food. As the availability of these compounds became limiting, there appeared autotrophic life forms in the sea with the capacity for photosynthesis. The appearance of these autotrophs (i.e. plants) facilitated the evolution of consumer heterotrophic organisms (i.e. animals). Furthermore, the evolution of oxygen as a by-product of photosynthesis resulted in the formation of the Ozone layer which shields this Earth from ultra-violet radiation. This radiation shield facilitated the colonization of shallow waters in the seas by both plants and animals, and subsequently the colonization of land.

In this process of evolution, most species have been doomed to loose, to be defeated and to become extinct because evolution itself is an ultimate existential game of groups and populations, in which nature is both the opponent and umpire selecting winners by rules that change unpredictably. Victory is thus a meagre privilege of staying for another round. The species on this planet today are thus only a small group of temporary winners. This process of progressive evolution resulted in the appearance of complex ecological systems in the tropics, and in the appearance of man on land about 3 million years ago. Man is presently the dominant species on this Earth.

Today, the marine environment accounts for 34% of the productivity of the biosphere, while the continental environment inclusive of freshwater systems 66%; and the continental environment accounts for 99.8% of the biomass with the tropical forests predominating (Whittaker and Likens 1973). Freshwater systems particularly rivers and swamps, contribute less than 5% to the continental biological productivity.

The net primary production of tropical drainage systems is low, being of the order of 10–34 grams of carbon/square metre/year in flowing waters (Ho and Furtado 1974; Lim and Furtado 1974; Likens 1973) due to swamp and riparian vegetation cover and to the abundance of allochthonous organic matter. However, inundated swamps with emergent vegetation have a net primary production of 3–8 tons/hectare/year (Ikusima 1974). As a consequence, the drainage waters are essentially heterotrophic below these macrophytes with the detritivore – decomposer food chains predominating. There is an abundance of aquatic insects, detritivorous fish, desmids and other organisms in these drainage systems (Ho and Furtado 1974; Johnson 1973). Several groups of aquatic insects and freshwater fishes have their centre of origin in tropical South East Asia, which is perhaps the only region with a long continuous history of tropical climates. By and large, however these taxa of organisms are poorly known, leaving much want for basic taxonomic studies which are critical for ecological and physiological work, and which are difficult because of intense speciation in this region.

When forests in river basins are manipulated for economic production, the exposure and enrichment of the drainage waters results in a change in their trophic dynamics from heterotrophy to autotrophy. The decomposer – detritivore food chain is disrupted resulting in the impairment of the capacity of the drainage system to process organic wastes, thus causing pollution (Vallentyne 1972, FAO 1969, Vollenweider 1968, WHO 1969). This capacity of the drainage system may also be impaired by other forms of pollution such as from mining and industrial activities (Ho 1973; Cheng 1977; CAP 1976). There are still very few detailed studies on the impact of landscape perturbations on the trophic dynamics of drainage waters in tropical South East Asia.

The drainage waters in tropical South East Asia are colonised by flora and fauna of both terrestrial and aquatic origin (Hynes 1970). By and large, the biology and ecology of these life forms is poorly understood. Nevertheless, these organisms may serve as good indicators of the health or trophic capacity of the drainage systems. In Malaysia, dominant plants (Corner 1940), plant species associations (Merton 1962), fish species associations (Tweedie 1961; Ali 1978; Johnson 1967b) and odonate species associations (Furtado 1969) have been used to classify different drainage water types. Mosquito species associations have been used to typify small containers (Hodgkin 1945; Colless 1956). Studies are in progress to examine the utility of other taxonomic groups of organisms such as mayflies (Lee Yoke Shum, per. comm.) and freshwater fish parasites (Lim and Furtado, in press). The constraints to such studies are taxonomic and physiological. A good indicator species must be integrative of the environmental conditions in an ecosystem, particularly the aquatic system. For example, the interrelationship of some Malaysian drainage waters is demonstrated by odonates which are the dominant predators among invertebrates (Furtado 1969):

(A) Piptospatha stream:

Non-torrential : Sg. Kongsi Lapan

Sundacypha petiolata Echo modesta

#### Torrential

Sg. Rumput Sg. Temson

## Devadatta a. argyoides

Euphaea o. ochracea Rhinocypha fenestrella Coeliccia albicauda Calicnemia chaseni Indocnemis orang

(B) Saraca Stream:

Upper

Sg. Gombak trib Sg. Ampang Euphaea o. ochracea

Rhinocypha perforata limbata Zygonyx iris malayana

Megalogomphus sumatranus

Viddle

Sg. Gombak

Sg. Tanglir

Neurobasis c. chinensis Gomphidia a. abbotti Onychothemis coccineaor O. culminicola

(C) Saraca stream:

Lower

Sg. Klang Sg. Chongkok Copera marginipes

Ictinogomphus decoratus melaenops Onychothemis t. testacea Libellago I. lineata

Open hill-country ponded stream: Ampang Reservoir

Sg. Klang (ponded)

(D) Open lowland stream: (Neram river)

Sg. Pelampas

Ictinogomphus decoratus melaenops Nannophya pygmaea

Macromia cincta Libellago I. lineata (E) Blackwater & forested lowland stream:

Sg. Lumut Sg. Bukit Cherakak Prodasineura interrupta

Libellago aurantiaca Elattaneura analis Euphaea impar

Forested hill-country ponded stream:

Sg. Kongsi Lapan (ponded) Copera v. vittata Tyriobapta toorida

(F) Lowland Ponded Streams:

Mining pool/Fish pond

Sg. pelumut (ponded)

Pandanus swamp

Sg. Ayer Hitam (ponded) C

A SHORE THE PERSON AND A SHORE AND A

Lacking unique species

Chalybeothemis fluviatilis Rhyothemis obsolescus

Pseudagrion coomansi

There are several points that emerge from this analysis:

- (1) Of the 77 species of odonates used in this classification of water systems, only some were of indicator value because of their unique and localised ecological and behavioural relationships to the water. Several species more could not be utilised in this analysis because of inadequate knowledge of their taxonomy, lifehistory and/or behaviour.
- (2) The ponding of a drainage water does not change its relationship to the running water very significantly, because many species inhabiting the pool regions of streams are able to colonise and adapt to the impoundment.
- (3) Under (E) there is an artificial relationship between blackwater forested lowland streams and a forested hill-country pondedstream, due probably to the abundance of common species with wide ecological ranges.

- (4) Similarly, under (F), there is an artificial relationship between lowland ponded – streams related to the Neram rivers and lowland Pandanus (Rassau) swamp and blackwater stream related to the Rassau rivers.
- (5) In the association analysis utilised, all species have equal weightage contributing thereby to these anomalous relationships between water bodies. There is a need therefore to develop methodologies in which certain species, such as rarer species, may be given greater importance value because of their vital nature to the dynamics of the ecological system.
- (6) There is a need to verify such association analysis with others, and with further biological particularly eco-physiological work.

Similarly, algal and acquatic plants (Arumugam 1976; Arumugam and Furtado, in press) and fishes (Yap 1978) have been used to characterise different stages in the eutrophication and recovery of a reservoir.

Besides aquatic organisms, certain terrestrial organisms may be used as indicators of the land-water interactions, especially organisms with a symbiotic organic relationship such as arboreal animals and primates, ant-plants, amphibians and mammals, insect pollinators (Appanah 1979), and soil fauna (Leow 1975). However, in most cases, the taxonomic constraints prevail.

### Water, Life and Man

Development or modernization is a continuous progressive process concerning the social evolution of man (Sauer 1971; Alatas 1972). The two critical constraints for this process becoming autonomous and indigenous are: (1) the rapid growth in scientific and technological capabilities of man to intervene in vital ecological processes, and (2) the lag in conceptual understanding of this socio-ecological transformation and of the psychosocial processes for managing change more effectively (McHale and Cordell 1974; Alatas 1971; Wilson 1971). The process of social transformation is the only problem common to both development and environmental conservation, and is rooted in a complex set of socio-cultural conditions in South East Asia (Furtado 1978). Emphasis on economic development is inducing rapid social changes and a variety of impacts on the trophic status of drainage waters:

- Effluent discharge of silt and toxic chemicals from mineral mining (Furtado 1978).
- (2) Soil erosion and surface runoff from land clearance for timber logging, fuelwood extraction, shifting cultivation, or for increasing agricultural lands (Burgess 1971; Singh 1972).
- (3) Nutrient and pesticide runoff from intensive agriculture especially wet ricefields (Farvar & Milton 1972).
- (4) Organic effluents from agro-industrial activities, and organic and inorganic effluents from other industries (Ho 1973; Bishop 1971 and 1973).
- (5) Impact of major engineering works such as highways, urban development, impoundments, and irrigation systems (Goodland and Irwin 1975; Ackermann, White and Worthington 1973).

Drainage water is thus affected by transportation or consumptive uses, more so by the latter, and by increasing scales of manipulation involving one, more than one or a continental system of river basins. Because of these multiple and often conflicting impacts such as Sungei Juru (CAP 1976) and forest catchments for watersupply, river basin authorities are urgently needed for managing effective water use and control, especially where none exist as in Malaysia.

It is evident from the foregoing that rivers are throughflow systems governed by higher units such as landscapes or biogeocoenoses (Sukachev 1964). They are analogous to the excretory system of the landscape, and thus are invaluable indicators of the development and conservation dynamics of the landscape. It is invaluable therefore to develop dynamic models, incorporating the concept of indicator species, for the estimation and prediction of water quality and quantity in relation to land use manipulation (Sandhu 1978; Whitehead and Young 1979).

Although Malaysia and several South East Asian nations possess a water surplus, already about 20% of runoff is used in various ways (Daniel and Cheong 1976) leaving only a margin of 10–15% for further development. These nations have thus reached a critical threshold where water is a decisive factor in economic development, especially

because of high consumptive rates in agriculture (10,000 tons of water/ ton of product) and industry (1,000 tons of water/ton of product). Indeed, as noted by E.A. Ackerman of the Carnegie Institute in 1959:

"The critical factor governing the development of life lies in the availability of sources of water".

These critical signs need to be considered seriously if we are to avoid the same forces that caused the decline of ancient civilizations such as in Mesopotamia, Nile and the Indus Valley, all of which thrived on irrigation systems.

The threat to future development in tropical South East Asia may indeed be an apparent and not a real one, to misunderstanding of the harsh sociobiological conditions surrounding modernisation, the flexible strategies needed for contemporary social change, and the interdependencies for organic growth (Janzen 1973; Furtado 1978). The future prospect for development relies heavily on conservation of resources, research and education for management of resources, and on planning and strategies for the development of resources. Planning for successful modernisation is in turn dependent on (1) a continual focus on the issues pertaining to total human and social development, (2) a continual incorporation of ecological perspectives into socio-economic development plans, and (3) a continual refinement of land capability classifications and regional development plans with information from various branches of research, particularly impacts of resource and social development, and ecological guidelines for development (Dasmann, Milton and Freeman 1973).

Strategies for modernisation need to be flexible for effective development, incorporating ingredients such as (1) the maintenance of a dynamic and sustainable relationship between man and his environment; (2) the nurturing of healthy competition at all levels to discover unique ways of modernisation at low social costs and risks (3) the utilization of a stochastic rather than a deterministic approach for optimising social change; (4) the formation of somewhat autonomous, small-scale, viable human communities for long-term stability; and (5) the incorporation of ecological principles in economic and political decision-making, particularly ecological resilience indicators for managing the unknown in the future (Dasmann, Milton and Freeman 1973; Omo-Fadaka 1975; Watt, Molloy, Varshney, Weeks and Wirosardjono 1977; Jacobs 1977; Furtado 1978). It is in this context that ecology, particularly of drainage systems and of indicator organisms, has an important role; and that biology can be in the forefront of development planning namely in the emerging discipline of landscape planning or ecology, or environmental design. Landscape ecology is not entirely a new science; the term has been used for budgets in Germany for centuries. It embodies the study of the dynamic factors impinging on resources (i.e. matter, energy, space, time and diversity) and the processes governing resource systems, the development and conservation of resources, spatial systems, and distributive systems of power and benefits.

I would like to conclude with this quote from Joseph W. Krutch, a naturalist:

"There is a paradox at the heart of the necessary 'struggle for existence', and the paradox is simply this: Neither man nor any other animal can afford to triumph in that struggle too completely. Unconditional surrender is a self-defeating formula — even in the war against insect pests. To the victor belong the spoils in nature also, but for a time only. Where there are no more spoils to be consumed, the victor dies."

#### Overview

In conclusion, it is evident that three aspects of zoology and ecology need academic emphasis:

- Taxonomy and ecology especially of animals in drainage waters, and with symbiotic relationships to land-water interactioon.
- (2) Ecological dynamics of land-water interactions particularly in relation to organic matter flows, perturbation impacts, and dynamic models for predicting and forecasting.
- (3) Biological indicators for development planning particularly those reflecting the trophic status and resilience of ecological systems.

Such emphasis would depend on inputs, support and efforts not only of materials (e.g. museum, equipment) but also of personnel.

#### References

- Ackermann, W.C., White, G.F. and Worthington, E.B., (eds). 1973. Man-made lakes: Their problems and environmental effects. American Geophysical Union, Washington, D.C.
- Alatas, S.H., 1972. Modernisations and social change: Studies in modernisation, religion, social change and development in South East Asia. Angus & Robertson Publ. Sydney.
- Appanah, S., 1979. The ecology of insect pollination of some tropical rain forest trees. 235 pp. Ph.D. Thesis, University of Malaya, Kuala Lumpur.
- Ali Awang, 1978. The feeding ecology of some riverine fishes from Sungei Langat at Ulu Langat, Selangor. 68 pp. B.Sc. Ecology Thesis, Department of Zoology, University of Malaya, Kuala Lumpur.
- Arumugam, P.T., 1976. Recovery of an eutrophicated reservoir, Subang Lake, Malaysia. 110 pp. M.Sc. Thesis, University of Malaya, Kuala Lumpur.
- Arumugam, P.T. and Furtado, J.I., in press. *Eutrophication of a Malay*sian reservoir: Effects of agro-industrial effluents. Proceedings of the Vth International Symposium of Tropical Ecology April 1979, Kuala Lumpur.
- Bishop, J.E., 1971. Limnology of a small Malayan river: Sungai Gombak. Ph.D. Thesis, University of Malaya, Kuala Lumpur.
- Bishop, J.E., 1973. Limnology of a small Malayan river: Sungai Gombak. Monogr. Biologicae. W. Junk, Hague.
- Burgess, P.F., 1971. The effect of logging on hill dipterocarp forests. Mal. Nat. J., 24, 231–237.
- CAP, 1976. Pollution: Kuala Juru's battle for survival. 73 pp. Consumers Association of Penang, Penang.
- Cheng, T.L., 1977. The Sungai Langat with special reference to pollution., B.Sc. Honours (Zoology) Thesis, Department of Zoology, University of Malaya, Kuala Lumpur.
- Chia, L.S., 1974. Seasonal rainfall distribution over West Malaysia. 33 pp. Paper presented at the IBP Synthesis Meeting, 12–18 August, 1974. Kuala Lumpur.
- Clapham, Jr., W.B., 1973. Natural ecosystems. Macmillan Co., New York.

- Colless, D.H., 1956. Studies in the systematics and biology of the culicine mosquitoes of Singapore. 230 pp. Ph.D. Thesis, University of Malaya (deposited at the Medical Library, University of Singapore).
- Corner, E.J.H., 1940. Wayside trees of Malaya, Vol. 1. 770 pp. Government Printers, Singapore.
- Dale, W.L., 1959. The rainfall of Malaya, Part 1. J. trop. Geogr., 13, 23-37.
- Daniel, J.G. and Cheong, C.L., 1976. Status of water resources activities and development in Malaysia. Report of the International Workshop on Energy, Resources and the Environment. February 1975, Malaysia, pp. 61–70. Malaysian Centre for the Development Studies, Kuala Lumpur.
- Dasmann, R.F., Milton, J.P. and Freeman, R.H., 1973. *Ecological prin*ciples for economic development. 252 pp. John Wiley, London.
- F.A.O., 1969. The influence of man on the hydrologic cycle. FAO/SC/ HYMIDEC/20 55 pp. U.N. Food & Agricultural Organization, Rome.
- Farvar, M.T. and Milton, J.P. (eds.), 1972. The careless technology: Ecology and international development. 1030 pp. Natural History Press, Garden City, N.Y.
- Fittkau, E.J., Irmler, U., Junk, W.J., Reiss, F. and Schmidt, G.W., 1975. Productivity, biomass, and population dynamics in Amazonian water bodies. pp. 289–312 in "Tropical ecological systems: Trends in terrestrial and aquatic research", eds. F.B. Golley & E. Medina. Springer Verlag, New York.
- Furtado, J.I. 1969. Ecology of Malaysian odonates: Biotope and association of species. Verh. Internat. Verein. Limnol., 17, 863–887.
- Furtado, J.I. 1978. The status and future of the tropical moist forest in the Southeast Asia. pp. 73–120 in "Developing economics in South East Asia and the environment" eds. C. MacAndrews and L.S. Chia. McGraw-Hill Singapore.
- Golley, F.B. McGinnis, J.T., Glements, R.G. Child, G.I. and Duever, M.J., 1975. *Mineral cycling in a tropical moist forest ecosystem*. University of Georgia Press, Athens.
- Goodland, R.J.A. and Irwin, H.S. 1975. *Amazon jungle: Green hell to red desert?* Landscape Planning, 1, 123–254.

- Gunaratnam, D.J. 1974. Monitoring changes to the hydrologic cycle due to land development by the use of computer models. 11 pp. Paper presented at the IBP Synthesis Meeting, August 1974, Kuala Lumpur.
- Ho, S.C. 1973. The ecology of a lowland stream: Sungei Renggam, with special reference to pollution. 170 pp. M.Sc Thesis, University of Malaya, Kuala Lumpur.
- Ho; S.C. and Furtado, J.I., 1974. Some aspects of the Limnology of lowland streams in West Malaysia. 21 pp. Paper presented at the IBP Synthesis Meeting, August 1974, Kuala Lumpur.
- Hodgkin, E.P., 1945. The Anopheles of Malaya. 48 pp. Perth, Western Australia.
- Hynes, H.B.N., 1970. *The ecology of running waters* University of Toronto Press, Toronto.
- Ikusima, I., 1974. Primary production of Tasek Bera with special reference to Lepironia swamps. 9 pp. Paper presented at the IBP Synthesis Meeting, August 1974, Kuala Lumpur.
- Jacob, F., 1977. Evolution and tinkering. Science, 196, 1161–1166.
- Janzen, D., 1973. Tropical agroecosystems. Science, 182, 1212-1219.
- Johnson, D.S., 1967a. On the chemistry of waters in Southern Malaya and Singapore. Arch. Hydrobiol, 63, 477–486.
- Johnson, D.S., 1967b. Distributional patterns in Malayan freshwater fish. Ecology, 48, 722-730.
- Johnson, D.S., 1973b. Freshwater life. pp. 103–127 inn "Animal life and nature in Singapore". ed. S.H. Chuang. Singapore University Press, Singapore.
- Kobayashi, J., 1959. Chemical investigations on river waters of southeastern Asiatic countries. Ber. Chara. Inst. Landwirtsch. Biologie, 11, 167–233.
- Leigh, C., 1974. Surface wash on two slopes in the Pasoh Forest Reserve, Negri Sembilan. 16 pp. Paper presented at the IBP Synthesis Meeting August 1974, Kuala Lumpur.
- Leigh, C., and Low, K.S., 1973. An appraisal of the flood situation in West Malaysia. pp. 57–72 in "Proceedings of the Symposium on Biological Resources and National Development" eds. E. Soepadmo and K.G. Singh. Malayan Nature Society, Kuala Lumpur.

Leow, K.C., 1975. *Meiofauna of decomposing leaves of some rainforest trees.* 78 pp. M.Sc. Thesis, University of Malaya, Kuala Lumpur.

- Likens, G.E., 1973. Primary production: Freshwater ecosystems, Human Ecology, 347–356.
- Lim, L.H. and Furtado, J.I., in press. *Freshwater fish host species and their parasites from Tasek Bera.* Proceedings of the Vth International Symposium of Tropical Ecology, April 1979, Kuala Lumpur.
- Lim, R.P. 1974. Limnological studies on a Malaysian freshwater swamp, Tasek Bera, Pahang. 114 pp. M.Sc. Thesis, University of Malaya, Kuala Lumpur.
- Lim, R.P. and Furtado, J.I., 1974. Primary production and plank ton at Tasek Bera, Malaysia, 3 pp. Paper presented at the IBP Synthesis Meeting August 1974, Kuala Lumpur.
- Low, K.S., 1972. Interception loss in the humid forested are as (with special reference to Sungei Lui catchment, West Malaysia). Malay. Nat. J. 25, 104–111.
- Manokaran, N., 1974. A preliminary study of the cycling of nutrients in various forms of water movement in the Pasoh Forest Ecosystem. 19 pp. Paper presented at the IBP Synthesis Meeting, August 1974, Kuala Lumpur.
- McHale, J. and Cordell, M., 1974. World trend and alternative futures Open Grants Paper No. 1. East-West Centre, Honolulu, Hawaii.
- Merton, F., 1962. A visit to the Tasek Bera. Malay. Nat. J., 16, 103-110.
- Mizonu, T. and Mori, S., 1969. Preliminary hydrobiological research of some South East Asian inland waters. Rep. Osaka Kyoiku Univ., 18, 119–165.
- Omo-Fadaka, J., 1975. Development. The Third way. Alternatives, 1, 23-53.
- Pereira, H.C., 1962. Hydrological effects of changes in land use in some East African catchments. East African Agric. For. J.; Spec. Iss., 27, 131 pp.
- Sandhu, H.S., 1978. Impact of landuse on water resources in the west coast of Peninsular Malaysia. 151 pp. M.Sc. Thesis, University of Malaya, Kuala Lumpur.
- Sandhu, H.S., Gunaratnam, D. and Furtado, J.I. in press. *The impact* of landuse on the quantity of runoff. Proceedings of the Vth International Symposium of Tropical Ecology, April 1979, Kuala Lumpur.

- Gauer, C.O., 1971. The agency of man on earth. pp. 51-71 in "Omega: Murder of the ecosystem and suicide of man". ed. P.K. Anderson, W.C. Brown & Co., Dubuque, Iowa.
- Singh, G., 1972. Report on watershed conditions in the lower Mekong basin and recommendations. A draft. 79 pp. Mekong Secretariat, UN.ESCAP, Bangkok.
- Sioli H., 1975a. Tropical rivers as expressions of their terrestrial environments. pp. 275–288 in "Tropical ecological systems: Trends in terrestrial and aquatic research". eds F.B. Golley and E. Medina. Springer Verlag, New York.
- Sioli H., 1975b. Amazon tributaries and drainage basins. pp. 199- in "Coupling of land and water systems". ed. A.D. Hasler. Springer Verlag, New York.
- Stewart, C.D., 1930. The rainfall of Malaya. Malayan Agric. J., 18, 530-540.
- Sukachev, V.N., 1964. Fundamentals of forest biogeocoenology. Oliver and Boyd, London.
- Swank, W.T. and Douglass, J.E. 1974. Streamflow greatly reduced by converting deciduous hardwood stands to pine. Science, 185, 857–859.
- Teoh, T.S., 1973. Some effects of Hevea plantations on rainfall redistribution. pp. 73-84. in "Proceedings of the Symposium on Biological Resources and National Development". Eds. E. Soepadmo and K.G. Singh. Malayan Nature Society, Kuala Lumpur.
- Tweedie, M.W.F., 1961. Malayan fish fauna and its future. pp. 230-231, in "Nature conservation in Western Malaysia" eds. J. Wyatt-Smith and P.R. Wycherley. Malayan Nature Society, Kuala Lumpur.
- Vallentyne, J.R., 1972. Freshwater supplies and pollution: Effects of the demophoric explosion on water and man. pp. 181-211 in "The environmental future" ed. N. Polunin. Macmillan Press, London.
- Verghese, S., 1975. Production and decomposition of swamp forest litter. 61 pp. B.Sc. Honours (Ecology) Thesis, Department of Zoology, University of Malaya, Kuala Lumpur.
- Verghese, S. and Furtado, J.I., in press. Litter production in a freshwater swamp forest, Tasek Bera, Malaysia. Proceedings of the Vth International Symposium of Tropical Ecology, April 1979, Kuala Lumpur.

- Vollenweider, R.A., 1968. Scientific fundamentals of the eutrophication of lakes and flowing waters, with particular reference to nitrogen and phosphorus as factors in eutrophication. DAS/CS1/68.27, 159 pp. Organization for Economic Cooperation & Development, Paris.
- Watt, K.E.F., Molloy, C.F., Varshney, C.K., Weeks, D. and Wirosardjono, S. 1977. *The unsteady state: Environmental problems,* growth, and culture. East-West Centre, Honolulu.
- W.H.O., 1969. *Water pollution in developing countries*. WHO Technical Rep. Ser. No. 404, 38 pp. World Health Organization, Geneva.
- Whitehead, P. and Young, P., 1979. *Water quality in river systems:* Monte-Carlo analysis. Water Resources Res., 15, 451–459.
- Whittaker, R.H., & Likens, G.E., 1973. *Primary production: The biosphere and Man.* Human Ecology, 1, 357–269.
- Wilson, D., 1971. Asia awakes: A continent in transition. Mentor Books, New York.
- Wycherley, P.R., 1967. *Rainfall probability tables for Malaysia*. Planting Manual No. 12, 85 pp. Rubber Research Institute of Malaysia, Kuala Lumpur.
- Yap, S.Y., 1978. The biology and ecology of Osteochilus hasselti (C. & V) with particular reference to its fisheries in Subang Reservoir, Malaysia. M.Sc. Thesis, University of Malaya, Kuala Lumpur.